

I'm pronounced "mol · lus · cum
con · ta · gi · o · sum"

APPOINTMENT GUIDE

Use this appointment guide before talking to your dermatologist about YCANTH[™], and treat molluscum right from the start.

As the name suggests, molluscum contagiosum, or molluscum, is a common, highly contagious skin disease caused by a poxvirus.

It spreads through skin-to-skin contact with someone with bumps and by sharing objects like:




Sports equipment



Dirty laundry



Towels



Toys



INDICATION:

YCANTH (cantharidin) topical solution, 0.7% is indicated for the topical treatment of molluscum contagiosum in adult and pediatric patients 2 years of age and older.

IMPORTANT SAFETY INFORMATION:

What warnings should I know about YCANTH?

- Do not get YCANTH in the mouth, nose, or eyes, and do not let YCANTH touch healthy skin. Life threatening or fatal toxicities can occur if YCANTH is taken by mouth. Avoid contact with areas of the body that have been treated, including contact by mouth. Damage to the eyes can occur if YCANTH comes in contact with the eyes. If YCANTH gets in the eyes, rinse eyes with water for at least 15 minutes.

Please see Important Safety Information throughout and click [here](#) for full Prescribing Information for YCANTH, including Patient Information.

DID YOU KNOW?

SCRATCHING CAN CAUSE THE BUMPS TO SPREAD

The bumps usually appear on children's arms, legs, and face, with a typical count of 10 to 30 bumps but as many as 100. Bumps can spread to other areas of the body or to family and friends.


Molluscum can be itchy, tender, or uncomfortable and have social impacts. Because of its contagious nature, it can prevent your child from being able to participate in their usual activities. The bumps can even lead to social isolation or bullying.

REFLECT AND SHARE

The bumps on my child's _____
bother them the most because they _____

Example: The bumps on my child's arms bother them the most because they itch and are seen by other kids.

MOLLUSCUM CAN LAST 5 YEARS



Left untreated, symptoms can last an average of 13 months but could persist for 5 years, and there is a risk of prolonged complications.

IMPORTANT SAFETY INFORMATION (cont'd):

What warnings should I know about YCANTH™? (cont'd)

- Local skin side effects at the application site may occur, including blistering, itching, pain, discoloration, and redness. Do not get YCANTH in the mouth, nose, or eyes, and do not let YCANTH touch healthy skin. If YCANTH contacts any unintended surface or healthy skin, immediately remove. If severe blistering, severe pain, or other severe skin side effects occur, wash off YCANTH immediately and contact your healthcare provider.

Please see Important Safety Information throughout and click [here](#) for full Prescribing Information for YCANTH, including Patient Information.

Ycanth™
(cantharidin) TOPICAL SOLUTION 0.7%



MOST PATIENTS OR PARENTS REPORT HOUSEHOLD SPREAD

No two families are exactly alike, so your reasons for treating molluscum are personal. Maybe you want to take back control of the infection, reduce the risk of spreading it to others, or prevent your child from spreading it to other parts of their body.


DID YOU KNOW?

REFLECT AND SHARE

I want to treat my child's molluscum _____

so we can _____

Example: I want to treat my child's molluscum before August so we can take care of it before swimming starts.



Y...canth you just wait and see?!

What if my doctor suggests that I wait and see?

YCANTH™ is a new molluscum treatment option for you and your dermatologist. Until now, there wasn't an FDA-approved treatment available, and the FDA has advised consumers against using unapproved products and treatments.

CONSIDER ASKING THEM:

- Q. If we start treatment with YCANTH today, when can we expect complete clearance of the bumps?
- Q. Has YCANTH helped your patients with molluscum return to their activities?
- Q. Are there any risks or benefits associated with YCANTH that we haven't discussed?

FOR AGES
2+

IMPORTANT SAFETY INFORMATION (cont'd):

What warnings should I know about YCANTH? (cont'd)


- YCANTH is flammable, even after drying. Do not expose YCANTH-treated areas to fire, flame, or smoke until YCANTH is washed off.
- Do not use YCANTH if you are allergic to any of the ingredients in YCANTH. The active ingredient in YCANTH is cantharidin. The inactive ingredients are acetone, camphor, castor oil, denatonium benzoate, ethanol, gentian violet, hydroxypropyl cellulose, and nitrocellulose.

Please see Important Safety Information throughout and click [here](#) for full Prescribing Information for YCANTH, including Patient Information.

Ycanth[™]
(cantharidin) TOPICAL SOLUTION 0.7%

DID YOU KNOW?

Time to pack my bags!



YCANTH™ CAN HELP START THE HEALING PROCESS AND CLEAR THE BUMPS IN JUST A FEW TREATMENTS

YCANTH PROVIDES RESULTS YOU CAN SEE

During your in-office visit, your dermatologist will apply a small, targeted amount of YCANTH directly to each bump. YCANTH causes small blisters on your skin to help remove the virus, so you can see it working.



Day 1

Before treatment



Day 2

After first treatment



12 weeks



End of treatment

Images taken from clinical trials. Photos are not representative of all patients.

The most common local skin side effects of YCANTH occurred at the application site and included blistering, itching, pain, discoloration, and redness. Avoid contact with areas of the body that have been treated, including contact by mouth. YCANTH is flammable, even after drying.

IMPORTANT SAFETY INFORMATION (cont'd):

What are the possible side effects of YCANTH?

The most common side effects are local skin side effects at the application site, including blistering, pain, itching, scabbing, redness, discoloration, dryness, swelling, and loss of the outer layer of skin at the application site. These local skin side effects are expected and are related to the anticipated blistering caused by YCANTH.

Please see Important Safety Information throughout and click [here](#) for full Prescribing Information for YCANTH, including Patient Information.

Ycanth™
(cantharidin) TOPICAL SOLUTION 0.7%

INDICATION:

YCANTH[™] (cantharidin) topical solution, 0.7% is indicated for the topical treatment of molluscum contagiosum in adult and pediatric patients 2 years of age and older.

IMPORTANT SAFETY INFORMATION:

What warnings should I know about YCANTH?

- Do not get YCANTH in the mouth, nose, or eyes, and do not let YCANTH touch healthy skin. Life threatening or fatal toxicities can occur if YCANTH is taken by mouth. Avoid contact with areas of the body that have been treated, including contact by mouth. Damage to the eyes can occur if YCANTH comes in contact with the eyes. If YCANTH gets in the eyes, rinse eyes with water for at least 15 minutes.
- Local skin side effects at the application site may occur, including blistering, itching, pain, discoloration, and redness. Do not get YCANTH in the mouth, nose, or eyes, and do not let YCANTH touch healthy skin. If YCANTH contacts any unintended surface or healthy skin, immediately remove. If severe blistering, severe pain, or other severe skin side effects occur, wash off YCANTH immediately and contact your healthcare provider.
- YCANTH is flammable, even after drying. Do not expose YCANTH-treated areas to fire, flame, or smoke until YCANTH is washed off.
- Do not use YCANTH if you are allergic to any of the ingredients in YCANTH. The active ingredient in YCANTH is cantharidin. The inactive ingredients are acetone, camphor, castor oil, denatonium benzoate, ethanol, gentian violet, hydroxypropyl cellulose, and nitrocellulose.

What are the possible side effects of YCANTH?

The most common side effects are local skin side effects at the application site, including blistering, pain, itching, scabbing, redness, discoloration, dryness, swelling, and loss of the outer layer of skin at the application site. These local skin side effects are expected and are related to the anticipated blistering caused by YCANTH.

Do other medications interact with YCANTH?

It is not currently known if any medications interact with YCANTH.

Can I use YCANTH if I'm pregnant?

The potential risk of YCANTH for major birth defects, miscarriage, or maternal or fetal side effects is unknown. Given that YCANTH is applied to the outside of the mother's skin, use is not expected to result in exposure of YCANTH to unborn babies.

Can I use YCANTH if I'm breastfeeding?

Do not use YCANTH on areas of the mother's body where YCANTH may come in contact with the breastfeeding child's mouth or eyes.

What should I do if YCANTH is swallowed?

Taking cantharidin by mouth has resulted in kidney failure, blistering and severe damage to the stomach and intestines, excessive bleeding or clotting, seizures, and weakness or paralysis. Seek medical attention immediately if YCANTH is accidentally ingested.

These are not all the possible side effects of YCANTH. Call your doctor for medical advice about side effects.

You are encouraged to report side effects to the FDA at 1-800-FDA-1088 or at www.fda.gov/medwatch. You may also report side effects to Verrica Pharmaceuticals Inc. at 1-877-VERRICA (1-877-837-7422). Local skin side effects are expected and should be reported if they are severe.

Please click [here](#) for full Prescribing Information for YCANTH, including Patient Information.